

# ON FRIENDLY INDEX SETS OF TREES

Ebrahim Salehi

Department of Mathematical Sciences

University of Nevada Las Vegas

Las Vegas, NV 89154-4020.

ebrahim.salehi@unlv.edu

Sin-Min Lee

Department of Computer Science

San Jose State University

San Jose, CA 95192.

lee@cs.sjsu.edu

ABSTRACT. For a graph  $G = (V, E)$  and a coloring  $f : V(G) \rightarrow \mathbb{Z}_2$  let  $v_f(i) = |f^{-1}(i)|$ .  $f$  is said to be friendly if  $|v_f(1) - v_f(0)| \leq 1$ . The coloring  $f : V(G) \rightarrow \mathbb{Z}_2$  induces an edge labeling  $f^* : E(G) \rightarrow \mathbb{Z}_2$  defined by  $f^*(xy) = f(x) + f(y) \forall xy \in E(G)$ , where the summation is done in  $\mathbb{Z}_2$ . Let  $e_f(i) = |f^{*-1}(i)|$ . The friendly index set of the graph  $G$ , denoted by  $FI(G)$ , is defined by

$$FI(G) = \{|e_f(1) - e_f(0)| : f \text{ is a friendly vertex labeling of } G\}.$$

In this paper we will determine the friendly index set of certain classes of trees, which in turn will verify the validity of the conjecture that the elements of friendly index set of any tree form an arithmetic progression.

**Key Words:** coloring, friendly index set, Fibonacci and Lucas trees.

**AMS Subject Classification:** 05C15, 05C25, 05C78

## 1. INTRODUCTION

Let  $G = (V, E)$  be a graph and  $f : V(G) \rightarrow \mathbb{Z}_2$  a vertex labeling (coloring) of  $G$ . For  $i \in \mathbb{Z}_2$ , let  $v_f(i) = |f^{-1}(i)|$ . The coloring  $f$  is said to be *friendly* if  $|v_f(1) - v_f(0)| \leq 1$ . For example, consider the graph  $G$ , illustrated in Figure 1, which has six vertices. To have a friendly coloring,

the condition  $|v_f(1) - v_f(0)| \leq 1$  implies that three vertices should be marked 0 and the other three 1.




FIGURE 1. A typical friendly labeling of  $G$ .

Any vertex labeling  $f : V(G) \rightarrow \mathbb{Z}_2$  induces an edge labeling  $f^* : E(G) \rightarrow \mathbb{Z}_2$  defined by  $f^*(xy) = f(x) + f(y) \forall xy \in E(G)$ . For  $i \in \mathbb{Z}_2$ , let  $e_f(i) = |f^{*-1}(i)|$ . The number  $N(f) = |e_f(1) - e_f(0)|$  is called the *friendly index* of  $f$ . The *friendly index set* of the graph  $G$ , denoted by  $FI(G)$ , is defined by

$$FI(G) = \{N(f) : f \text{ is a friendly coloring of } G \}.$$

A friendly coloring  $f : V(G) \rightarrow \mathbb{Z}_2$  is called *maximum friendly coloring* of  $G$  if its friendly index is the maximum element of  $FI(G)$ . Note that if  $f : V(G) \rightarrow \mathbb{Z}_2$  is a friendly coloring, so is its inverse coloring  $g : V(G) \rightarrow \mathbb{Z}_2$  defined by  $g(v) = 1 - f(v) \forall v \in V(G)$ . Moreover,  $N(g) = N(f)$ . The induced edge labeling of the graph  $G$ , Figure 1, is illustrated in Figure 2. It is easy to see that  $FI(G) = \{0, 2, 6\}$ .




FIGURE 2. The induced edge labeling of  $G$ .

Three friendly colorings of  $G$  that provide these numbers are presented in Figure 3.

In 1978, Ibrahim Cahit [2, 3, 4] introduced the concept of cordial labeling as weakened version of the less tractable graceful and harmonious labeling. A graph  $G$  is said to be cordial if it admits a friendly labeling with index 0 or 1. Cahit, among other facts, proved that

1. every tree is cordial;
2. The complete graph  $K_n$  is cordial if and only if  $n \leq 3$ ;
3. The complete bipartite graph  $K(m, n)$  is cordial ( $m, n \in \mathbb{N}$ );
4. The wheel  $W_n$  is cordial if and only if  $n \not\equiv 3 \pmod{4}$ ;

5. In an Eulerian graph  $G = (p, q)$  if  $p \equiv 0 \pmod{4}$ , then it is not cordial.

M. Hovay [8], later generalized the concept of cordial graphs and introduced  $A$ -cordial labelings. A graph  $G$  is said to be  $A$ -cordial if it admits a labeling  $f : V(G) \rightarrow A$  such that for every  $i, j \in A$ ,

$$|v_f(i) - v_f(j)| \leq 1 \text{ and } |e_f(i) - e_f(j)| \leq 1.$$

Cordial graphs have been studied extensively. Interested readers are referred to a number of relevant literature that are mentioned in the bibliography section, including [1, 5, 7, 9, 11, 13, 17]. In this paper, we will focus on group  $A = \mathbb{Z}_2$ . The *friendly index set* of a graph  $G$ , denoted by  $FI(G)$ , is defined to be  $\{N(f) = |e_f(1) - e_f(0)| : f \text{ is friendly coloring}\}$ . When there is no ambiguity we will drop the subscript  $f$ . Note that if 0 or 1 is in  $FI(G)$ , then  $G$  is cordial. Thus the concept of friendly index sets could be viewed as a generalization of cordiality.




FIGURE 3.  $FI(G) = \{0, 2, 6\}$ .

In general, the elements of  $FI(G)$  do not necessarily form an arithmetic progression. However, Lee and Ng [15] conjectured that the elements of the friendly index set of any tree will form an arithmetic progression. This conjecture has been verified for several classes of trees. To further verify the validity of this conjecture, in this paper, we will find the friendly index sets of several classes of trees. First a few well-known results [14]:

**Theorem 1.1.** For any graph  $G$  with  $q$  edges,  $FI(G) \subset \{q - 2i : i = 0, 1, 2, \dots, \lfloor q/2 \rfloor\}$ .

**Theorem 1.2.** Let  $1 \leq m \leq n$ . For the complete bipartite graph  $K_{m,n}$  we have

$$FI(K_{m,n}) = \begin{cases} \{(m - 2i)^2 : 0 \leq i \leq \lfloor m/2 \rfloor\} & \text{if } m + n \text{ is even;} \\ \{i(i + 1) : 0 \leq i \leq m\} & \text{if } m + n \text{ is odd.} \end{cases}$$

For any  $n \geq 1$ , the complete bipartite graph  $K(1, n)$  is called a *star* and is denoted by  $ST(n)$ . Stars are the only trees of diameter 2, for which we have:

**Corollary 1.3.**  $FI(ST(n)) = \begin{cases} \{0, 2\} & \text{if } n \text{ is even;} \\ \{1\} & \text{if } n \text{ is odd.} \end{cases}$

**Theorem 1.4.** *The friendly index set of a full binary tree with depth  $d > 1$  is  $\{0, 2, 4, \dots, 2^{d+1} - 4\}$ .*

## 2. TREES WITH PERFECT MATCHING

**Definition 2.1.** A *matching* in a graph is a set of edges with no shared endpoints. A matching  $M$  in a graph  $G$  is said to be a *perfect matching* if every vertex of  $G$  is incident with an edge in  $M$ .

Note that every graph with perfect matching has even number of vertices. Also, stars  $ST(n)$  ( $n \geq 2$ ) are among trees that do not have a perfect matching. As a result, if we identify one of the end-vertices of  $ST(n)$  ( $n \geq 3$ ) with a vertex of any graph  $G$ , the resulting graph does not have a perfect matching.




FIGURE 4. An example of a tree with perfect matching.

**Observation 2.2.** *Every terminal edge of a tree with perfect matching  $M$  is in  $M$ .*

**Observation 2.3.** *A tree  $T$  ( $|T| \geq 3$ ) with perfect matching has at least one  $P_3$ , the path of order 3, pendant. In fact, the two end portion of the longest path of  $T$  would have  $P_3$  pendants.*

**Theorem 2.4.** *If  $T = (p, q)$  is a tree with perfect matching, then  $FI(T) = \{1, 3, 5, \dots, q\}$ .*

*Proof.* Let  $M$  be a perfect matching of the tree  $T$ . We will proceed by induction on  $|M|$ . Clearly, the theorem is true for  $|M| = 1, 2$ . Assume the theorem is true whenever the the perfect matching has  $n$  elements. Let  $T$  be a tree with a perfect matching  $M$  such that  $|M| = n + 1$ . This implies

that  $|E(T)| = 2|M| + 1$ . By observation 2.3,  $T$  has at least a  $P_3$  pendant. Let  $u \sim v \sim w$  be such a pendant with  $vw$  being the terminal edge. Consider  $T' = T - \{v, w\}$ , which is a tree with perfect matching. In fact,  $M - vw$  is a perfect matching in  $T'$  with  $|M - vw| = n$ . By the induction hypothesis  $FI(T') = \{1, 3, \dots, 2n - 1\}$ .

Let  $f : V(T') \rightarrow \{0, 1\}$  be an arbitrary friendly coloring of  $T'$  and define  $\phi : V(T) \rightarrow \{0, 1\}$  by

$$\phi(x) = \begin{cases} f(x) & \text{if } x \neq v, w; \\ f(u) & \text{if } x = v; \\ 1 - f(u) & \text{if } x = w. \end{cases}$$

Then  $\phi$  is a friendly coloring of  $T$  with  $e_\phi(1) = 1 + e_f(1)$  and  $e_\phi(0) = 1 + e_f(0)$ . Therefore,

$$|e_\phi(1) - e_\phi(0)| = |e_f(1) - e_f(0)|,$$

which implies that  $FI(T') = \{1, 3, \dots, 2n - 1\} \subset FI(T)$ .

Now, let  $g : V(T') \rightarrow \{0, 1\}$  be a maximum friendly coloring of  $T'$  with  $e_g(1) = 2n - 1$  (or  $e_g(0) = 0$ ) and define  $\psi : V(T) \rightarrow \{0, 1\}$  by

$$\psi(x) = \begin{cases} f(x) & \text{if } x \neq v, w; \\ 1 - f(u) & \text{if } x = v; \\ f(u) & \text{if } x = w. \end{cases}$$

Then  $\psi$  is a friendly coloring of  $T$  with  $e_\psi(1) = 2 + e_g(1) = 2n + 1$ , which implies that  $2n + 1 \in FI(T)$ .  $\square$

With a similar argument presented in the proof of theorem 2.4 one can show that

**Theorem 2.5.**  $FI(P_n) = \{n - 1 - 2i : 0 \leq i \leq \lfloor \frac{n-1}{2} \rfloor\}$ , where  $P_n$  ( $n \geq 2$ ) is the path of order  $n$ .

The theorem 2.5 implies that for any natural number  $n$  there is a connected graph  $G$  such that  $FI(G) = \{n - 2i : i = 1, 2, \dots, \lfloor n/2 \rfloor\}$ , which is an arithmetic progression with common difference being 2.

**Problem 2.6.** *Given any even integer  $d > 2$ , find a connected graph whose friendly index set forms an arithmetic progression with common difference  $d$ .*

Frucht and Harary [6] introduce the corona of two graphs  $G$  and  $H$ , denoted by  $G \odot H$ , to be the graph with base  $G$  such that each vertex  $v \in V(G)$  is joined to all vertices of a separate copy of  $H$ .

**Corollary 2.7.** *Given a tree  $T$  with  $q$  edges, the coronation  $T \odot K_1$  is a tree with perfect matching and its friendly index set is  $\{1, 3, 5, \dots, 2q + 1\}$ .*




FIGURE 5.  $FI(P_n \odot K_1) = \{1, 3, \dots, 2n - 1\}$ .

### 3. FIBONACCI AND LUCAS TREES

Fibonacci Trees, denoted by  $FT_n$ , are defined inductively as follows:  $FT_1$  is the trivial tree with one vertex,  $FT_2$  is the path  $P_2$ , and for  $n \geq 3$ ,  $FT_n = (V_n, E_n)$  is the binary tree of root  $r_n$ , whose left and right children are  $FT_{n-1}$  and  $FT_{n-2}$ , respectively.





FIGURE 6. Graphs of the first five Fibonacci trees.

**Observation 3.1.** *Let  $p_n = |V_n|$ , be the number of vertices of  $FT_n$ . Then  $p_n = A_{n+2} - 1$ , where  $A_n$  is the  $n^{\text{th}}$  Fibonacci number.*

**Theorem 3.2.** For  $n \geq 3$ , the friendly index set of  $FT_n$  is  $\{|E_n| - 2i : i = 0, 1, 2, \dots, \lfloor q_n/2 \rfloor\}$ . Moreover, every element of the index set can be obtained by a friendly coloring  $f : V_n \rightarrow \mathbb{Z}_2$  with  $N(f) = e_f(1) - e_f(0)$ , and the color of root is 1.

*Proof.* We proceed by induction on  $n$ . Clearly the theorem is true for  $n = 2, 3, 4$ . For  $FT_4$ , it is illustrated in Figure 7. Suppose it is true for all  $k$  with  $3 \leq k \leq n$  and consider  $FT_{n+1}$ . Let  $b_1 \in \{|E_n| - 2i : i = 1, 2, \dots, \lfloor q_n/2 \rfloor\}$  and  $b_2 \in \{|E_{n-1}| - 2i : i = 1, 2, \dots, \lfloor q_{n-1}/2 \rfloor\}$ . Also, let  $\phi : V_n \rightarrow \mathbb{Z}_2$  be a friendly coloring of  $FT_n$  with  $e_\phi(1) - e_\phi(0) = b_1$  and  $\psi : V_{n-1} \rightarrow \mathbb{Z}_2$  be a friendly coloring of  $FT_{n-1}$  with  $e_\psi(1) - e_\psi(0) = b_2$ . Without loss of generality, by considering the inverse coloring if necessary, we may assume that the roots of  $FT_n$  and  $FT_{n-1}$  are labeled 1 or 0.




FIGURE 7. Four desired friendly colorings of  $FT_4$ .

Define the function  $f : V_{n+1} \rightarrow \mathbb{Z}_2$  by

$$f(x) = \begin{cases} \phi(x) & \text{if } x \in V_n; \\ \psi(x) & \text{if } x \in V_{n-1}; \\ j & \text{if } x = r_{n+1}, \end{cases}$$

where  $j \in \mathbb{Z}_2$ . To make sure that  $f$  is a friendly coloring we will consider two cases:

**Case A.**  $|V_n|$  and  $|V_{n-1}|$  are even. By defining  $f(r_{n+1}) = 1$ ,  $f$  would be a friendly coloring of  $FT_{n+1}$ . If the roots of  $FT_n$  and  $FT_{n-1}$  are labeled 0, then  $N(f) = b_1 + b_2 + 2$ . If the roots of  $FT_n$  and  $FT_{n-1}$  are labeled differently, then  $N(f) = b_1 + b_2$ . Hence,  $b_1 + b_2$ ,  $b_1 + b_2 + 2$  are in the friendly index set of  $FT_{n+1}$  and these numbers will completely determine the set.

**Case B.** One of the numbers  $|V_n|$  or  $|V_{n-1}|$  is odd and the other one is even. Assume that  $|V_n|$  is odd,  $v_\phi(1) = v_\phi(0) + 1$ , and the root of  $FT_n$  is labeled 1. This implies that  $f(r_{n+1}) = 0$ . If

the root of  $FT_{n-1}$  is labeled 1, then  $N(f) = b_1 + b_2 + 2$ . If the root of  $FT_{n-1}$  is labeled 0, then  $N(f) = b_1 + b_2$ . Hence,  $b_1 + b_2$ ,  $b_1 + b_2 + 2$  are in the friendly index set of  $FT_{n+1}$  and these numbers will completely determine this set. Finally, we consider the inverse coloring of  $f$ , if needed, to guarantee that  $f(r_{n+1}) = 1$ .  $\square$

**Definition 3.3.** *Lucas Trees, denoted by  $LT_n$ , are defined inductively as follows:  $LT_1$  is the trivial tree with one vertex,  $LT_2$  is the path  $P_3$  with the middle vertex being its root, and for  $n \geq 3$ ,  $LT_n = (V_n, E_n)$  is the binary tree of root  $r_n$ , whose left and right children are  $LT_{n-1}$  and  $LT_{n-2}$ , respectively.*




FIGURE 8. Graphs of the first five Lucas trees.

**Observation 3.4.** *Let  $p_n = |V_n|$ , be the number of vertices of  $LT_n$ . Then  $p_n = 2A_{n+1} - 1$ , where  $A_n$  is the  $n^{th}$  Fibonacci number.*

**Theorem 3.5.** *For  $n \geq 3$ , the friendly index set of the Lucas tree  $LT_n = (V_n, E_n)$  is  $\{0, 2, 4, \dots, |E_n|\}$ . Moreover, every element  $b$  of this index set can be obtained by a friendly coloring  $f : V_n \rightarrow \mathbb{Z}_2$  with the following properties:*

- (a)  $f(r_n) = 1$ ;
- (b)  $b = N(f) = e_f(1) - e_f(0)$ ;

- (c) If  $b \equiv 0 \pmod{4}$ , then  $v_f(1) = v_f(0) + 1$ ; and
- (d) If  $b \equiv 2 \pmod{4}$ , then  $v_f(0) = v_f(1) + 1$ .

*Proof.* We proceed by induction on  $n$ . Clearly the theorem is true for  $n = 2, 3, 4$ . For  $LT_4$ , it is illustrated in Figure 9. Suppose it is true for all  $k$  with  $3 \leq k \leq n$  and consider  $LT_{n+1}$ . Let  $b_1 \in \{0, 2, 4, \dots, |E_n|\}$  and  $b_2 \in \{0, 2, 4, \dots, |E_{n-1}|\}$ . We wish to show that  $b_1 + b_2$  and  $b_1 + b_2 + 2$  belong to the index set of  $LT_{n+1}$ .




FIGURE 9. Five desired friendly colorings of  $LT_4$ .

Using the induction hypothesis, let  $\phi : V_n \rightarrow \mathbb{Z}_2$  be a friendly coloring of  $FT_n$  with  $b_1 = e_\phi(1) - e_\phi(0)$  and  $\psi : V_{n-1} \rightarrow \mathbb{Z}_2$  be a friendly coloring of  $FT_{n-1}$  with  $b_2 = e_\psi(1) - e_\psi(0)$ . Also, let  $\phi(r_n) = \psi(r_{n-1}) = 1$ .

First, we wish to prove that  $0 \in FI(LT_{n+1})$ . Choose  $b_1 = 0$  and  $b_2 = 2$ . Then  $v_\phi(1) = v_\phi(0) + 1$ , and  $v_\psi(0) = v_\psi(1) + 1$ . This situation is summarized in Figure 10.




FIGURE 10.  $+i$  refers to extra  $i$  in the coloring of the graph.

Define the function  $f : V_{n+1} \rightarrow \mathbb{Z}_2$  by

$$f(x) = \begin{cases} \phi(x) & \text{if } x \in V_n; \\ \psi(x) & \text{if } x \in V_{n-1}; \\ 1 & \text{if } x = r_{n+1}, \end{cases}$$

which is a friendly coloring of  $LT_{n+1}$ , with  $N(f) = b_1 + b_2 - 2 = 0$ .

Next, assume  $b_1$  or  $b_2 \neq 0$ ; for example let  $b_1 \neq 0$ . Also, let  $\lambda : V_n \rightarrow \mathbb{Z}_2$  be a friendly coloring of  $FT_n$  with  $b_1 - 2 = e_\lambda(1) - e_\lambda(0)$ . Depending on whether  $b_1$  or  $b_2$  are multiple of 4 or not, we will consider the following two cases:

**Case 1.** At least one of  $b_1$  or  $b_2$  is a multiple of 4; assume  $b_2 \equiv 0 \pmod{4}$ . In this case  $v_\psi(1) = v_\psi(0) + 1$ . The function  $g : V_{n+1} \rightarrow \mathbb{Z}_2$  defined by

$$g(x) = \begin{cases} \phi(x) & \text{if } x \in V_n; \\ \psi(x) & \text{if } x \in V_{n-1}; \\ 0 & \text{if } x = r_{n+1}, \end{cases}$$

is a friendly coloring of  $LT_{n+1}$ , with  $N(g) = b_1 + b_2 + 2$ . Also, the function  $h : V_{n+1} \rightarrow \mathbb{Z}_2$  defined by

$$h(x) = \begin{cases} \lambda(x) & \text{if } x \in V_n; \\ \psi(x) & \text{if } x \in V_{n-1}; \\ 0 & \text{if } x = r_{n+1}, \end{cases}$$

is a friendly coloring of  $LT_{n+1}$ , with  $N(h) = b_1 + b_2$ . The inverse coloring of  $g, h$  will satisfy the conditions of the theorem.




FIGURE 11.  $b_2 \equiv 0 \pmod{4}$ .

**Case 2.**  $b_1, b_2 \equiv 2 \pmod{4}$ . In this case  $b_1 - 2$  is multiple of 4 and the function  $h : V_{n+1} \rightarrow \mathbb{Z}_2$ , defined above, is a friendly coloring of  $LT_{n+1}$ , with  $N(h) = b_1 + b_2$ . Again, the inverse coloring of  $h$ , will satisfy the conditions of the theorem.

This proves that if  $b_1$  or  $b_2 \neq 0$ , then  $b_1 + b_2$  and  $b_1 + b_2 + 2$  are members of  $FI(LT_{n+1})$ . □

We conclude the paper with the improved version of the conjecture:

**Conjecture 3.6.** *The elements of the friendly index set of any tree form an arithmetic progression with common difference 2.*

#### REFERENCES

- [1] M. Benson and S-M. Lee, On Cordialness of Regular Windmill Graphs, *Congressus Numerantium* **68** (1989), 49-58.
- [2] I. Cahit, Cordial Graphs: a weaker version of graceful and harmonious graphs, *Ars Combinatoria* **23** (1987), 201-207.
- [3] I. Cahit, On Cordial and 3-equitable Graphs, *Utilitas Mathematica* **37** (1990), 189-198.
- [4] I. Cahit, Recent Results and Open Problems on Cordial Graphs, *Contemporary Methods in Graph Theory*, Bibliographisches Inst. Mannheim (1990), 209-230.
- [5] N. Cairnie and K. Edwards, The Computational Complexity of Cordial and Equitable Labellings, *Discrete Mathematics* **216** (2000), 29-34.
- [6] R. Frucht and F. Harary, On the Corona of Two Graphs, *Aequationes Mathematicae* **4** (1970), 322-325.
- [7] Y.S. Ho, S-M. Lee, and S.C. Shee, Cordial Labellings of the Cartesian Product and Composition of Graphs, *Ars Combinatoria* **29** (1990), 169-180.
- [8] M. Hovay, A-cordial Graphs, *Discrete Mathematics* **93** (1991), 183-194.
- [9] W.W. Kirchherr, On the Cordiality of Certain Specific Graphs, *Ars Combinatoria* **31** (1991), 127-138.
- [10] W.W. Kirchherr, NEPS Operations on Cordial Graphs, *Discrete Mathematics* **115** (1993), 201-209.
- [11] S. Kuo, G.J. Chang, and Y.H.H. Kwong, Cordial Labeling of  $mKn$ , *Discrete Mathematics* **169** (1997) 121-131.
- [12] Y.H. Lee, H.M. Lee, and G.J. Chang, Cordial Labelings of Graphs, *Chinese J. Math.* **20** (1992), 263-273.
- [13] S-M. Lee and A. Liu, A Construction of Cordial Graphs from Smaller Cordial Graphs, *Ars Combinatoria* **32** (1991), 209-214.
- [14] S-M. Lee and H.K. Ng, On Friendly Index Sets of Bipartite Graphs, to appear in *Ars Combin.*
- [15] S-M. Lee and H.K. Ng, On Friendly Index Sets of Spiders, manuscript.
- [16] E. Seah, On the Construction of Cordial Graphs, *Ars Combinatoria* **31** (1991), 249-254.
- [17] M.A. Seoud and A.E.I. Abdel Maqsood, On Cordial and Balanced Labelings of Graphs, *J. Egyptian Math. Soc.* **7** (1999), 127-135.
- [18] S.C. Shee and Y.S. Ho, The Cordiality of one-point Union of  $n$  Copies of a Graph, *Discrete Mathematics* **117** (1993), 225-243.
- [19] S.C. Shee and Y.S. Ho, The Cordiality of the Path-union of  $n$  Copies of a Graph, *Discrete Mathematics* **151** (1996), 221-229.